



Start the conversation with your doctor



Did you know your diabetes can change as your body changes over time? If you've been taking oral diabetes treatments or injectable therapy and you're still struggling to lower your A1c, it may be time to talk to your doctor. Here are a few questions you can ask to find out if SOLIQUA 100/33 may be right for you.

Questions to ask:


- 1 What is my A1c now, and what A1c should I be aiming for?
- 2 I diet, exercise and take my diabetes medicine. Why isn't that enough to lower my A1c?
- 3 My A1c is still too high. Is this a sign that my diabetes treatment plan may need to change? Are there other signs I should watch for?
- 4 Why might a treatment like SOLIQUA 100/33—which combines a long-acting insulin with a non-insulin diabetes medicine—be the right next step for me?

Please see additional questions on page 2



Sign up for the \$9* CO-PAY!

Eligible commercially insured patients save on SOLIQUA 100/33: Pay as little as \$9* for a 30-day supply. Terms & conditions apply. Apply online today at: soliqua100-33.com/savings



*This offer is for commercially insured patients and is not valid for prescriptions covered by or submitted for reimbursement under Medicare, Medicaid, VA, DOD, TRICARE, or similar federal or state programs, including any state pharmaceutical programs. Void where prohibited by law. Savings card carries maximum savings of \$365 per pack, up to 2 packs for each 30-day supply, for the duration of the program. Savings may vary depending on patient's out-of-pocket costs. Upon registration, patient receives all program details. Sanofi US reserves the right to rescind, revoke, or amend the program without notice.

SOLIQUA 100/33 is an injectable prescription medicine that contains 2 diabetes medicines, insulin glargine and lixisenatide, that may improve blood sugar [glucose] control in adults with type 2 diabetes, when used with diet and exercise.

- It has not been studied in people with a history of pancreatitis.
- It is not recommended for people who also take lixisenatide or other medicines called GLP-1 receptor agonists.
- It is not for use in people with type 1 diabetes, diabetic ketoacidosis.
- It has not been studied in people who have a stomach problem that causes slow emptying [gastroparesis] and is not for people with slow emptying of the stomach.
- It has not been studied in people who also take a short-acting [prandial] insulin.
- **It is not known if SOLIQUA 100/33 is safe and effective in children under 18 years of age.**

Important Safety Information

What is the most important information I should know about SOLIQUA 100/33?

Do not share your SOLIQUA 100/33 pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

Please see full Important Safety Information for SOLIQUA 100/33 on pages 3 and 4.

Please see full Prescribing Information on the soliqua100-33.com website where you printed this form.



Start the conversation with your doctor

Questions to ask (continued):

- 5 Is SOLQUA 100/33 right for me? Are there side effects I should be aware of?
- 6 If you prescribe SOLQUA 100/33, what should I know about the pen? Will my dose change over time?
- 7 If SOLQUA 100/33 is right for me, when would I administer it and what will my routine be like?

Important Safety Information (Continued)

SOLQUA 100/33 can cause serious side effects, including inflammation of the pancreas, which may be severe and lead to death.

Before using SOLQUA 100/33, tell your doctor if you have had pancreatitis, stones in your gallbladder (cholelithiasis), or a history of alcoholism. These medical problems may make you more likely to get pancreatitis.

Stop taking SOLQUA 100/33 and call your healthcare provider right away if you have pain in your stomach area (abdomen) that is severe, and will not go away. The pain may be felt in the back area. The pain may happen with or without vomiting.

Who should not use SOLQUA 100/33?

Do not use SOLQUA 100/33 if you:

- are having an episode of low blood sugar (hypoglycemia)
- are allergic to insulin glargine, lixisenatide, or any of the ingredients in SOLQUA 100/33. Symptoms of a severe allergic reaction with SOLQUA 100/33 may include swelling of the face, lips, tongue, or throat, fainting or feeling dizzy, problems breathing or swallowing, very rapid heartbeat, severe rash or itching, or low blood pressure.

Before using SOLQUA 100/33, tell your healthcare provider about all your medical conditions, including if you:

- have or have had problems with your pancreas, your kidneys, or your liver, stones in your gallbladder, or a history of alcoholism.
- have heart failure or other heart problems. If you have heart failure, it may get worse while you take thiazolidinediones (TZDs).
- have severe problems with your stomach, such as slowed emptying of your stomach or problems digesting food.
- are taking certain medicines called glucagon-like peptide 1 receptor agonists (GLP-1 receptor agonists).
- have had an allergic reaction to a GLP-1 receptor agonist.

Please see full Important Safety Information for SOLQUA 100/33 on pages 3-4.

Please see full Prescribing Information on the soliqua100-33.com website where you printed this form.



Start the conversation with your doctor

SOLIQUEA[®] 100/33
insulin glargine & lixisenatide injection
100 Units/mL & 33 mcg/mL

Important Safety Information

What is the most important information I should know about SOLIQUEA 100/33?

Do not share your SOLIQUEA 100/33 pen with other people, even if the needle has been changed. You may give other people a serious infection, or get a serious infection from them.

SOLIQUEA 100/33 can cause serious side effects, including inflammation of the pancreas, which may be severe and lead to death.

Before using SOLIQUEA 100/33, tell your doctor if you have had pancreatitis, stones in your gallbladder (cholelithiasis), or a history of alcoholism. These medical problems may make you more likely to get pancreatitis.

Stop taking SOLIQUEA 100/33 and call your healthcare provider right away if you have pain in your stomach area (abdomen) that is severe, and will not go away. The pain may be felt in the back area. The pain may happen with or without vomiting.

Who should not use SOLIQUEA 100/33?

Do not use SOLIQUEA 100/33 if you:

- are having an episode of low blood sugar (hypoglycemia)
- are allergic to insulin glargine, lixisenatide, or any of the ingredients in SOLIQUEA 100/33. Symptoms of a severe allergic reaction with SOLIQUEA 100/33 may include swelling of the face, lips, tongue, or throat, fainting or feeling dizzy, problems breathing or swallowing, very rapid heartbeat, severe rash or itching, or low blood pressure.

Before using SOLIQUEA 100/33, tell your healthcare provider about all your medical conditions, including if you:

- have or have had problems with your pancreas, your kidneys, or your liver, stones in your gallbladder, or a history of alcoholism.
- have heart failure or other heart problems. If you have heart failure, it may get worse while you take thiazolidinediones (TZDs).
- have severe problems with your stomach, such as slowed emptying of your stomach or problems digesting food.
- are taking certain medicines called glucagon-like peptide 1 receptor agonists (GLP-1 receptor agonists).
- have had an allergic reaction to a GLP-1 receptor agonist.
- are pregnant or breastfeeding or plan to become pregnant or to breastfeed. It is not known if SOLIQUEA 100/33 will harm your unborn baby or pass into your breast milk.

Tell your healthcare provider about all the medicines you take, including all prescription and over-the-counter medicines, vitamins, and herbal supplements. SOLIQUEA 100/33 may affect the way some medicines work. Before using SOLIQUEA 100/33, talk to your healthcare provider about low blood sugar and how to manage it.

Please see additional Important Safety Information for SOLIQUEA 100/33 on page 4.

Please see full Prescribing Information on the soliqua100-33.com website where you printed this form.



Start the conversation with your doctor

SOLQUA[®] 100/33
insulin glargine & lixisenatide injection
100 Units/mL & 33 mcg/mL

Important Safety Information (Continued)

How should I use SOLQUA 100/33?

- Do not change your dose without first talking to your healthcare provider.
- Check the pen label each time you inject to make sure you are using the correct medicine.
- **Do not take more than 60 units of SOLQUA 100/33 each day.** Do not take SOLQUA 100/33 with other GLP-1 receptor agonists.
- Only use SOLQUA 100/33 that is clear and colorless to almost colorless. If you see small particles, return it to your pharmacy for replacement.
- **Change (rotate) your injection sites within the area you chose with each dose** to reduce your risk of getting lipodystrophy (pitted or thickened skin) and localized cutaneous amyloidosis (skin with lumps) at the injection sites. **Do not** use the same spot for each injection or inject where the skin is pitted, thickened, lumpy, tender, bruised, scaly, hard, scarred or damaged.
- **Do not** remove SOLQUA 100/33 from the pen with a syringe.
- **Do not re-use or share needles with other people. You may give other people a serious infection, or get a serious infection from them.**
- **Check your blood sugar levels.** Ask your healthcare provider what your blood sugar should be and when you should check.

What are the possible side effects of SOLQUA 100/33?

SOLQUA 100/33 may cause serious side effects, including:

- **Serious allergic reactions.** Stop taking SOLQUA 100/33 and get help right away if you have any symptoms of a serious allergic reaction, including swelling of your face, lips, tongue, or throat, problems breathing or swallowing, severe rash or itching, fainting or feeling dizzy, and very rapid heartbeat.
- **Low blood sugar (hypoglycemia). Your risk for getting low blood sugar is higher if you take another medicine that can cause low blood sugar.** Signs and symptoms of low blood sugar may include headache, dizziness, drowsiness, sweating, weakness, irritability, hunger, blurred vision, fast heartbeat, feeling jittery, confusion, and anxiety.
- **Kidney problems (kidney failure).** In people who have kidney problems, diarrhea, nausea, and vomiting may cause a loss of fluids (dehydration), which may worsen kidney problems.
- **Low potassium in your blood (hypokalemia).**
- **Heart failure.** Taking certain diabetes pills called TZDs (thiazolidinediones) with SOLQUA 100/33 may cause heart failure in some people. This can happen even if you have never had heart failure or heart problems before. If you already have heart failure, it may get worse while you take TZDs with SOLQUA 100/33. Tell your healthcare provider if you have any new or worse symptoms of heart failure, including shortness of breath, swelling of your ankles or feet, or sudden weight gain. Treatment with TZDs and SOLQUA 100/33 may need to be adjusted or stopped if you have new or worse heart failure.

The most common side effects of SOLQUA 100/33 include low blood sugar (hypoglycemia), nausea, diarrhea, upper respiratory infection, stuffy or runny nose, and headache. Nausea and diarrhea usually happen more often when you first start using SOLQUA 100/33.

Please see full Prescribing Information on the soliqua100-33.com website where you printed this form.